

Deformation of the Dirac Equation

Mir Faizal¹ and Sergey I. Kruglov²

¹Department of Physics and Astronomy,
University of Waterloo, Waterloo,
Ontario N2L 3G1, Canada

²Department of Chemical and Physical Sciences,
University of Toronto, 3359 Mississauga Rd. North,
Mississauga, Ontario, L5L 1C6, Canada

Abstract

In this paper, we will first clarify the physical meaning of having a minimum measurable time. Then we will combine the deformation of the Dirac equation due to the existence of minimum measurable length and time scales with its deformation due to the doubly special relativity. We will also analyse this deformed Dirac equation in curved spacetime, and observe that this deformation of the Dirac equation also leads to a non-trivial modification of general relativity. Finally, we will analyse the stochastic quantization of this deformed Dirac equation on curved spacetime.

1 Introduction

All most all approaches to quantum gravity predict the existence of a minimum measurable length scale. For example, in string theory the smallest probe that can be used for analysing spacetime is the fundamental string. So, it is not possible to probe spacetime at length scales smaller than the string length scale. Thus, string theory comes naturally equipped with a minimum length scale, which is the string length scale [1]-[5]. A minimum length scale also exists in loop quantum gravity. In fact, this minimum length scale in loop quantum gravity turns the big bang into a big bounce [6]. The existence of a minimum length scales can also be inferred from black hole physics [7]-[8]. This is because the energy required to probe spacetime at length scales smaller than Planck length is more than the energy required to form a black hole in that region of spacetime. Thus, it can be inferred from black hole physics that any theory of quantum gravity should naturally come equipped with an minimum measurable length scale of the order of Planck scale.

The existence of a minimum length is not consistent with the usual Heisenberg uncertainty principle. This is because according to the usual Heisenberg uncertainty principle, it is possible to measure length with arbitrary accuracy if momentum is not measured. So, the usual Heisenberg uncertainty principle has to be generalized to generalized uncertainty principle, to make the existence of minimum length consistent with quantum mechanics [9]-[14]. However, the Heisenberg uncertainty principle is closely related to the Heisenberg algebra,

and so, the deformation of the usual Heisenberg uncertainty principle, will also lead to a deformation of the Heisenberg algebra [15]-[19]. This deformation of the Heisenberg algebra changes the momentum operator in the coordinate representation, and this will in turn produce correction terms for all quantum mechanical systems [20]-[23].

A different form of deformation of the Heisenberg algebra occurs due a modification of the special relativity to doubly special relativity [24]-[26]. Apart from the velocity of light, the Planck energy is also a universal constant in doubly special relativity. The doubly special relativity is motivated by a modification of the usual energy-momentum dispersion relation, which in turn is motivated by studies done on discrete spacetime [27], the spontaneous symmetry breaking of Lorentz invariance in string field theory [28], spacetime foam models [29], spin-network in loop quantum gravity [30], non-commutative geometry [31], and Horava-Lifshitz gravity [32]. In fact, it has also been possible to generalize doubly special relativity to doubly general relativity, and the resultant theory is known as gravity's rainbow [33]-[34]. The deformations of the Heisenberg algebra caused by the existence of a minimum measurable length and doubly special relativity have been combined into a single deformation of the Heisenberg algebra [35]-[37]. Corrections to various physical systems have been studied using this deformed Heisenberg algebra. For example, the transition rate of ultra cold neutrons in gravitational field received corrections due to this deformed Heisenberg algebra [38]. Similarly, the correction terms for the Lamb shift and the Landau levels have also been calculated [39]. One of the most interesting consequence of this deformed algebra is that it leads to a discretization of space [35]. This deformation of the Heisenberg algebra changes the form of the first quantized field theories, and second quantization of these deformed field theory has also been studied [40]-[43]. The deformation of the Wheeler-DeWitt equation corresponding to this algebra has been performed [44]. It was observed that due to such a deformation the big bang singularity can be avoided. This happens because there is a minimum value for the scale factor of the universe. So, the universe can not become smaller than this minimum value, and hence, the big bang singularity is avoided.

It may be noted that it is possible to define a system with minimum measurable time, and if the principle of covariance in relativity is taken seriously, then a minimum measurable length scale should also correspond to a minimum measurable time scale. This can be done by defining the observable time as the time of occurrence of an certain type of event associated with the appearance of some specified value of a dynamical quantity [45]-[50]. In fact, there are good reasons to treat the space and time on equal footing, as it has been argued that by treating the space and time on equal footing the black-hole information paradox gets naturally resolved [51]-[52]. Thus, it is possible to generalize the work on deformation of Heisenberg algebra based on minimum length scale to both minimum length and time scales. It may also be noted that another interesting deformation of quantum mechanics comes from stochastic quantization [53]-[56]. In fact, it has been argued that the Gell-Mann-Oakes-Renner relation has an analogous meaning in QCD, and this leads to the existence of a Thouless energy in QCD [57]-[59]. It has been possible to calculate this Thouless energy in QCD in the framework of stochastic quantization [60]. General relativity can be treated as gauge theory [61]-[63], and so, it is possible to apply the results obtained for stochastic quantization of gauge theories to gravity.

In this paper, we will first clarify what it means to have a minimum measurable time in a theory. This will be done by defining an minimum measurable value to the certainty with which the occurrence of an event of a certain type can be measured by the appearance of some specified value of a dynamical quantity. Then we will combine the deformation of the Heisenberg algebra caused by the doubly special relativity, with its deformation caused by existence of minimal measurable length and time scales. We will also complete this algebra by including a covariant deformation of its temporal part. The Dirac equation consistent with this deformed algebra Heisenberg will contain non-local fractional derivative terms. However, such terms will be given a formal meaning in the framework of harmonic extension of functions. We will also analyse the effect this deformation of Dirac equation has on general relativity. We observe that apart from deforming the matter action, this deformation also leads to a non-trivial modification of the gravitational part of the action. As general relativity can be viewed as a gauge theory, we will analyse the deformed Dirac equation on curved spacetime in the framework of stochastic quantization. We will perform our analysis semi-classically taking the gravitational field as a fixed background. We will also demonstrate that this stochastic quantization of the deformed Dirac equation can be done in the superspace formalism.

2 Deformation of Heisenberg Algebra

In this section, we first generalize the deformation of quantum mechanics caused by the existence of a minimum measurable length scale to the deformation caused by the existence of both minimum measurable length and time scales. The existence of minimum length causes the following deformation of the uncertainty principle [15]-[19],

$$\Delta x \Delta p = \frac{\hbar}{2} [1 + \beta (\Delta p)^2], \quad (1)$$

and this in turn deforms the Heisenberg algebra to

$$[x^i, p_j] = i\hbar [\delta_j^i + \beta p^2 \delta_j^i + 2\beta p^i p_j], \quad (2)$$

where $\beta = \beta_0 \ell_{Pl}^2 / \hbar^2$, β_0 is a constant normally assumed to be of order one, and $\ell_{Pl} \approx 10^{-35}$ m. The coordinate representation of the momentum operator can now be written as

$$p_i = i\hbar \partial_i (1 - \beta \hbar^2 \partial^i \partial_i). \quad (3)$$

Now a deformation of the Heisenberg algebra corresponds to the existence of a minimum measurable time is also expected to take place due to principles of covariance in relativity. Thus, we expect the uncertainty between energy and time to get deformed to

$$\Delta E \Delta t = \frac{\hbar}{2} [1 + \beta (\Delta E)^2], \quad (4)$$

where we have set $t \rightarrow it$. It may be noted that such a deformation has already been studied [46]. This deformation will in turn changes the representation of the Hamiltonian operator to

$$H = i\hbar \partial_t (1 - \beta \hbar^2 \partial_t^2). \quad (5)$$

It may be noted that it is possible to define time as an observable in quantum mechanics, with reference to the evolution of some non-stationary quantity [45]-[50]. Thus, the observable time is the time of occurrence of an event of a certain type, defined by the appearance of some specified value of a dynamical quantity. So, by viewing time as an observable in quantum mechanics, it is possible to give physical meaning to the commutator $[H, t] = i\hbar$. Now the deformation of the Hamiltonian operator caused by the existence of a minimum measurable time scale corresponds to the deformation of the commutator to $[H, t] = i\hbar[1+3\beta H^2]$, where again we have set $t \rightarrow it$. This would physically correspond to the case where there is an minimum measurable value to the certainty with which the occurrence of an event of a certain type can be measured by the appearance of some specified value of a dynamical quantity.

Now we will analyse the deformation of the Heisenberg algebra which incorporates both the deformations corresponding to the existence of minimum length and doubly special relativity. The deformation of the Heisenberg algebra caused doubly special relativity is given by

$$[x^i, p_j] = i\hbar[1 - \tilde{\beta}|p^k p_k|^{1/2}\delta_j^i + \tilde{\beta}^2 p^i p_j], \quad (6)$$

where $\tilde{\beta} = \ell_{Pl} \approx 10^{-35} m$. It is possible to combine the deformation occurring due to the existence of a minimum length scale and the deformation occurring due to doubly special relativity into a single deformation of the Heisenberg algebra [35]-[37]

$$\begin{aligned} [x^i, p_j] = & i\hbar \left[\delta_j^i - \alpha|p^k p_k|^{1/2}\delta_j^i + \alpha|p^k p_k|^{-1/2}p^i p_j \right. \\ & \left. + \alpha^2 p^k p_k \delta_j^i + 3\alpha^2 p^i p_j \right], \end{aligned} \quad (7)$$

where M_{Pl} is the Planck mass, $\alpha = \alpha_0/M_{Pl}c = \alpha_0\ell_{Pl}/\hbar$, and $M_{Pl}c^2 \approx 10^{19} GeV$ is the Planck energy. It has been suggested that the parameter α_0 could be situated at an intermediate length scale between Planck scale and electroweak scale [39]. If that is the case, then it will be possible to devise experiments to analyse the phenomenological effects of quantum gravitational phenomena. In fact, in that case, it might be possible to observe the upper bound for α_0 by using quantum optics techniques [64]. The coordinate representation of the momentum operator can now be written as

$$p_i = -i\hbar \left(1 - \hbar\alpha\sqrt{-\partial^j \partial_j} - 2\hbar^2\alpha^2\partial^j \partial_j \right) \partial_i. \quad (8)$$

Motivated by the deformation of the momentum operator corresponding to the existence of a minimal measurable time, we assume the following form for the deformed four dimensional momentum operator in the coordinate representation,

$$p_\mu = -i\hbar \left(1 - \hbar\alpha\sqrt{-\partial^\nu \partial_\nu} - 2\hbar^2\alpha^2\partial^\nu \partial_\nu \right) \partial_\mu. \quad (9)$$

This deformed momentum operator satisfies the following algebra,

$$\begin{aligned} [x^\mu, p_\nu] = & i\hbar \left[\delta_\nu^\mu - \alpha|p^\rho p_\rho|^{1/2}\delta_\nu^\mu + \alpha|p^\rho p_\rho|^{-1/2}p^\mu p_\nu \right. \\ & \left. + \alpha^2 p^\rho p_\rho \delta_\nu^\mu + 3\alpha^2 p^\mu p_\nu \right]. \end{aligned} \quad (10)$$

where again we have set $t \rightarrow it$. Here again time has to be viewed as an observable, for the temporal part of this commutator to be physically meaningful. This is again done by first defining the observable time as the time of occurrence of an event of a certain type at which some specified value gets assigned to a dynamical quantity [45]-[50]. Then viewing this algebra as the existence of a minimum measurable value to the certainty with which the occurrence of that event can be measured by the appearance of this specified value for that dynamical quantity. It may be noted that if time is taken as an observable, then motivated by the principle of covariance of space and time, it was expected that this deformed algebra should also included its temporal part. We will assume this to be a general form for the momentum operator in all field theories. Thus, we will be able to analyse the deformation of the Dirac equation by using this deformed momentum operator. We will also chose natural unites and set $c = \hbar = 1$.

3 Deformed Dirac Equation

In the previous section, we have analysed the deformation of the Heisenberg algebra caused the doubly special relativity along with the existence of minimum measurable length and time scales. In this section, we will analyse the Dirac equation corresponding to this deformation of the Heisenberg algebra. Now we can write the Dirac equation corresponding to this deformed Heisenberg algebra as follows [40]-[43],

$$i\gamma^\mu \left(1 - \alpha\sqrt{-\partial^\nu\partial_\nu} - 2\alpha^2\partial^\nu\partial_\nu\right) \partial_\mu\psi(x) + m\psi(x) = 0. \quad (11)$$

where γ^μ are Dirac's matrices obeying $\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2\eta^{\mu\nu}$, and $\psi(x)$ is bispinor [74].

This Dirac equation contains non-local fractional derivative term, however, this term can be given a formal meaning in the framework of harmonic extensions of a function [65]-[69]. We start by defining a harmonic extension of $\psi : R^n \rightarrow R$, which is a harmonic function $u : R^n \times (0, \infty) \rightarrow R$. The restriction of u to R^n coincides with $\psi : R^n \rightarrow R$. We can also define ∂_{n+1}^2 as the Laplacian in R^{n+1} , such that $x \in R^n$ and $y \in R$. Now for a given ψ , we can solve the Dirichlet problem given by $u(x, 0) = \psi(x)$ and $\partial_{n+1}^2 u(x, y) = 0$, to find u . Thus, for a given smooth function $C_0^\infty(R^n)$, there is a unique harmonic extension $u \in C^\infty(R^n \times (0, \infty))$.

A formal meaning can now be given to $\sqrt{-\partial^\mu\partial_\mu}$ by analysing its action on $\psi : R^n \rightarrow R$. We require in that case the harmonic extension of the ψ to satisfy $u : R^n \times (0, \infty) \rightarrow R$,

$$\sqrt{-\partial^\mu\partial_\mu}\psi(x) = -\frac{\partial u(x, y)}{\partial y} \Big|_{y=0}. \quad (12)$$

Here $u_y(x, y)$ is the harmonic extension of $\sqrt{-\partial^\mu\partial_\mu}\psi(x)$ to $R^n \times (0, \infty)$, as $u : R^n \times (0, \infty) \rightarrow R$ is a harmonic extension of $\psi : R^n \rightarrow R$. The repeated application of $\sqrt{-\partial^\mu\partial_\mu}$ on ψ can be written as

$$\begin{aligned} \left[\sqrt{-\partial^\mu\partial_\mu}\right]^2 \psi(x) &= \frac{\partial^2 u(x, y)}{\partial y^2} \Big|_{y=0} \\ &= -\partial_n^2 u(x, y) \Big|_{y=0}. \end{aligned} \quad (13)$$

This equation gives a formal meaning to $\sqrt{-\partial^\mu \partial_\mu}$. The harmonic extension of $\partial_\mu \psi(x)$ can be written as $\partial_\mu u(x, y)$ because $u(x, y)$ is the harmonic extension of the $\psi(x)$, and $\sqrt{-\partial^\mu \partial_\mu} \psi = -u_y(x, 0)$. Now as $u \in C^2(R \times (0, \infty))$, so, we can write

$$\sqrt{-\partial^\mu \partial_\mu} \partial_\nu \psi(x) = -\partial_\nu u_y(x, y) |_{y=0}. \quad (14)$$

Thus, any derivative commutes with this non-local operator $\sqrt{-\partial^\mu \partial_\mu}$,

$$\sqrt{-\partial^\mu \partial_\mu} \partial_\nu \psi(x) = \partial_\nu \sqrt{-\partial^\mu \partial_\mu} \psi(x). \quad (15)$$

Now we have given a formal meaning to the non-local Dirac equation with fractional derivative terms. We can also write this deformed Dirac equation in a matrix valued formalism. To do that we introduce the following functions,

$$\begin{aligned} \chi(x) &= -\frac{1}{m} \partial_5 \psi(x), \\ \phi(x) &= -\frac{1}{m^2} \partial^\mu \partial_\mu \psi(x) \\ &= -\frac{1}{m} \partial_5 \chi, \end{aligned} \quad (16)$$

where $\partial_5 \equiv \sqrt{-\partial^\mu \partial_\mu}$. Then this deformed Dirac equation can be written as

$$\begin{aligned} i\gamma^\mu (\partial_\mu \psi(x) + \alpha m \partial_\mu \chi(x) + 2\alpha^2 m^2 \partial_\mu \phi(x)) + m\psi(x) &= 0, \\ \partial_5 \psi(x) + m\chi(x) &= 0, \\ \partial_5 \chi(x) + m\phi(x) &= 0. \end{aligned} \quad (17)$$

We introduce a 12-component wave function, $\Psi(x) = (\psi(x), \chi(x), \phi(x))$, and write the deformed Dirac equation as

$$i\Gamma^\mu \partial_\mu \Psi(x) + i\Gamma^5 \partial_5 \Psi(x) + m\Psi(x) = 0, \quad (18)$$

where

$$\begin{aligned} \Gamma^\mu &= \gamma^\mu \otimes A, & \Gamma^5 &= I_4 \otimes B, \\ A &= \varepsilon^{1,1} + \alpha m \varepsilon^{1,2} + 2\alpha^2 m^2 \varepsilon^{1,3}, & B &= \varepsilon^{2,1} + \varepsilon^{3,2}. \end{aligned} \quad (19)$$

Here A is a projection matrix $A^2 = A$, $B^3 = 0$, and I_4 is a unit 4×4 -matrix. Thus, advantage of using this formalism is that the non-local fractional derivative term can be effectively written as a local term. Thus, using the notations $\Gamma^A = (\Gamma^\mu, \Gamma^5)$, and $\partial_A = (\partial_\mu, \partial_5)$, we can now write the deformed Dirac equation as

$$(i\Gamma^A \partial_A + m) \Psi(x) = 0, \quad (20)$$

where $A = 1, 2, 3, 4, 5$. It may be noted that this is effectively written as a local Dirac equation. Now in the momentum space we have the following result

$$(\mathcal{P} - m) \Psi(p) = 0, \quad (21)$$

where $\mathcal{P} = \Gamma^\mu p_\mu + \Gamma^5 p_5$, and so,

$$\mathcal{P} = \begin{pmatrix} \gamma^\mu p_\mu & \alpha m \gamma^\mu p_\mu & 2\alpha^2 m^2 \gamma^\mu p_\mu \\ p_5 I_4 & 0 & 0 \\ 0 & p_5 I_4 & 0 \end{pmatrix}. \quad (22)$$

We also have $A^2 = A$, $B^2 = \varepsilon^{3,1}$, $B^3 = 0$, and

$$\begin{aligned}\Gamma^\mu \Gamma^\nu + \Gamma^\nu \Gamma^\mu &= 2\eta^{\mu\nu} I_4 \otimes A, \\ \Gamma^\rho (\Gamma^\mu \Gamma^\nu + \Gamma^\nu \Gamma^\mu) &= 2\eta^{\mu\nu} \Gamma^\rho.\end{aligned}\quad (23)$$

4 Deformation of General Relativity

It is possible to write the deformation of any gauge theory by converting the deformed derivatives into deformed covariant derivatives [41]-[42]. In fact, for a Lie algebra $[T_A, T_B] = if_{AB}^C T_C$, the gauge field $A_\mu = A_\mu^A T_A$ can be used to write a covariant derivative $D_\mu = \partial_\mu + iA_\mu$. It has been demonstrated that it is possible to analyse a deformation of a gauge theory by deforming $-i(1 - \alpha\sqrt{-\partial^\nu \partial_\nu} - 2\alpha^2 \partial^\nu \partial_\nu) \partial_\mu$ to $-i(1 - \alpha\sqrt{-D^\nu D_\nu} - 2\alpha^2 D^\nu D_\nu) D_\mu$ [43]. In this paper, we analyse such a deformation of general relativity. It is possible to formulate the general relativity as a gauge theory by considering the translation group as a gauge group [61]-[62]. Even though the torsion plays a central role in this formalism, it can be demonstrated that this formalism is equivalent to general relativity. It is also possible to formulate the general relativity as a gauge theory by using the Lorentz group as a gauge group [63]. We expect that the gauge theories to get modified due to the deformation of the Heisenberg algebra, so, the deformation of the Heisenberg algebra will also lead to a non-trivial modification of general relativity. In this paper, we will analyse general relativity as a gauge theory whose gauge group is the Lorentz group. Now under the Lorentz transformation given by $x^a \rightarrow x^b \Lambda_b^a$, a fermion will transform as

$$\psi \rightarrow U\psi = \exp\left(\frac{i}{2}\Lambda^{ab}(x)\Sigma_{ab}\right), \quad (24)$$

where Σ_{ab} describes the generators of Lorentz group represented within Dirac spinor space as $\Sigma_{ab} = -i[\gamma_a, \gamma_b]/4$. Now we define $g_{\mu\nu}e_a^\mu e_b^\nu = \eta_{ab}$, and so, we can write a spinor covariant derivative as

$$D_a = e_a^\mu (\partial_\mu I + \frac{i}{2}\omega_\mu^{ab}\Sigma_{ab}) = e_a^\mu D_\mu. \quad (25)$$

Here ω_μ^{ab} is the spin connection which is given by

$$\begin{aligned}\omega_\mu^{ab} &= 2e^{\nu a}\partial_\mu e_\nu^b - 2e^{\nu b}\partial_\mu e_\nu^a - 2e^{\nu a}\partial_\nu e_\mu^b + 2e^{\nu b}\partial_\nu e_\mu^a \\ &\quad + e_{\mu c}e^{\nu a}e^{\rho b}\partial_\rho e_\nu^a - e_{\mu c}e^{\nu a}e^{\rho b}\partial_\nu e_\rho^c.\end{aligned}\quad (26)$$

The commutator of these two covariant derivative can be written as

$$[D_\mu, D_\nu] = A_{\mu\nu}^c D_c + \frac{i}{2}R_{\mu\nu}^{ab}\Sigma_{ab}. \quad (27)$$

We can set the torsion to zero, $A_{\mu\nu}^c = 0$, and write the Riemann tensor with two Minkowski indices as

$$R_{\mu\nu}^{ab} = \partial_\mu \omega_\nu^{ab} - \partial_\nu \omega_\mu^{ab} + \omega_\mu^{ac}\omega_{\nu c}^b - \omega_\nu^{ac}\omega_{\mu c}^b. \quad (28)$$

Here the spin connection ω_μ^{ab} transforms as

$$\omega_\mu^{ab} \rightarrow [U\omega_\mu U^{-1} - (\partial_\mu U)U^{-1}]^{ab}, \quad (29)$$

so, $R_{\mu\nu}^{ab}$ transforms as

$$R_{\mu\nu}^{ab} \rightarrow [U]_c^a R_{\mu\nu}^{cd} [U^{-1}]_d^b. \quad (30)$$

Now we can define the scalar curvature as $R = e_a^\mu e_b^\mu R_{\mu\nu}^{ab}$, and write the action, with $\kappa = 8\pi G$, as

$$S = \frac{1}{2\kappa} \int d^4x e R + \int d^4x e \bar{\psi} (i\gamma^a e_a^\mu D_\mu - m) \psi. \quad (31)$$

So, the Einsteins equation can now be written as

$$R_\mu^a - \frac{1}{2} e_\mu^a R = \kappa T_\mu^a, \quad (32)$$

where T_μ^a is the energy momentum tensor corresponding to the Dirac equations.

Now we can write the deformed Dirac equation in curved spacetime by using the assumption that the form of the coordinate representation of the deformed momentum operator will be the same in all field theories. So, we can replace the derivatives in the coordinate representation of the deformed momentum operator by covariant derivatives. Thus, the deformed Dirac equation can be written as [40]-[43]

$$\gamma^\mu \left(1 - \alpha \sqrt{-D^\nu D_\nu} - 2\alpha^2 D^\nu D_\nu \right) D_\mu \psi(x) + m\psi(x) = 0. \quad (33)$$

Now if we define $\mathcal{D}_\mu = (1 - \alpha \sqrt{-D^\tau D_\tau} - 2\alpha^2 D^\tau D_\tau) D_\mu$, then we can write the deformed commutator as

$$\begin{aligned} [\mathcal{D}_\mu, \mathcal{D}_\nu] &= \mathcal{A}_{\mu\nu}^c D_c + \frac{i}{2} \mathcal{R}_{\mu\nu}^{ab} \Sigma_{ab} \\ &= A_{\mu\nu}^c D_c + \frac{i}{2} R_{\mu\nu}^{ab} \Sigma_{ab} + \alpha \left[A(\alpha)_{\mu\nu}^c D_c + \frac{i}{2} R(\alpha)_{\mu\nu}^{ab} \Sigma_{ab} \right] \\ &\quad + \alpha^2 \left[A(\alpha^2)_{\mu\nu}^c D_c + \frac{i}{2} R(\alpha^2)_{\mu\nu}^{ab} \Sigma_{ab} \right]. \end{aligned} \quad (34)$$

So, we can write

$$\begin{aligned} \mathcal{A}_{\mu\nu}^c &= A_{\mu\nu}^c + \alpha A(\alpha)_{\mu\nu}^c + \alpha^2 A(\alpha^2)_{\mu\nu}^c, \\ \mathcal{R}_{\mu\nu}^{ab} &= R_{\mu\nu}^{ab} + \alpha R(\alpha)_{\mu\nu}^{ab} + \alpha^2 R(\alpha^2)_{\mu\nu}^{ab}, \end{aligned} \quad (35)$$

where $A(\alpha)_{\mu\nu}^c D_c + iR(\alpha)_{\mu\nu}^{ab} \Sigma_{ab}/2$ and $A(\alpha^2)_{\mu\nu}^c D_c + iR(\alpha^2)_{\mu\nu}^{ab} \Sigma_{ab}/2$ are correction terms generated from the deformation of Heisenberg algebra,

$$\begin{aligned} A_{\mu\nu}^c D_c + R_{\mu\nu}^{ab} \Sigma_{ab} &= [D_\mu, D_\nu], \\ A(\alpha)_{\mu\nu}^c D_c + R(\alpha)_{\mu\nu}^{ab} \Sigma_{ab} &= - \left[\sqrt{-D^\tau D_\tau} D_\mu, D_\nu \right] - \left[D_\mu, \sqrt{-D^\tau D_\tau} D_\nu \right], \\ A(\alpha^2)_{\mu\nu}^c D_c + R(\alpha^2)_{\mu\nu}^{ab} \Sigma_{ab} &= -2 [D^\tau D_\tau D_\mu, D_\nu] - 2 [D_\mu, D^\tau D_\tau D_\nu] \\ &\quad + \left[\sqrt{-D^\tau D_\tau} D_\mu, \sqrt{-D^\rho D_\rho} D_\nu \right]. \end{aligned} \quad (36)$$

We again set the torsion equal to zero, $A_{\mu\nu}^c = A(\alpha)_{\mu\nu}^c = A(\alpha^2)_{\mu\nu}^c = 0$, and write the deformed Riemann tensor with two Minkowski indices as

$$\mathcal{R}_{\mu\nu}^{ab} = \partial_\mu \omega_\nu^{ab} - \partial_\nu \omega_\mu^{ab} + \omega_\mu^{ac} \omega_{\nu c}^b - \omega_\nu^{ac} \omega_{\mu c}^b + R(\alpha)_{\mu\nu}^{ab} + R(\alpha^2)_{\mu\nu}^{ab} \quad (37)$$

Now as $\mathcal{D}_\mu = (1 - \alpha\sqrt{-\bar{D}^\nu D_\nu} - 2\alpha^2 D^\nu D_\nu)D_\mu$ also transforms as D_μ , the transformation $\mathcal{R}_{\mu\nu}^{ab}$ can be written as

$$\begin{aligned}\mathcal{R}_{\mu\nu}^{ab} &\rightarrow [U]_c^a \mathcal{R}_{\mu\nu}^{cd} [U^{-1}]_d^b \\ &= [U]_c^a (R_{\mu\nu}^{cd} + R(\alpha)_{\mu\nu}^{cd} + R(\alpha^2)_{\mu\nu}^{cd}) [U^{-1}]_d^b.\end{aligned}\quad (38)$$

Now we can define the deformed scalar curvature as $\mathcal{R} = e_a^\mu e_b^\mu \mathcal{R}_{\mu\nu}^{ab}$, and write the deformed action as

$$S = \frac{1}{2\kappa} \int d^4x e \mathcal{R} + \int d^4x e \bar{\psi} (i\gamma^a e_a^\mu D_\mu - m) \psi. \quad (39)$$

The deformed Einstein equations can also be written as

$$\mathcal{R}_\mu^a - \frac{1}{2} e_\mu^a \mathcal{R} = \kappa \mathcal{T}_\mu^a, \quad (40)$$

where $\mathcal{R}_\mu^a = \mathcal{R}_{\mu\nu}^{ab} e_\nu^b$ is the Ricci tensor generated from the deformed Riemann tensor, and $\mathcal{T}_\mu^a = T_\mu^a + \alpha T(\alpha)_\mu^a + \alpha^2 T(\alpha^2)_\mu^a$ is the deformed energy momentum tensor. It may be noted that apart from the deformation of the energy-momentum tensor corresponding to the deformation of the original theory, the gravitational action for this theory also gets deformed.

5 Stochastic Quantization

In this section, we will analyse the stochastic quantization of Dirac equation in curved spacetime. We will analyse this model semi-classically and treat gravity as a fixed classical background and analyse the stochastic quantization of Dirac equation on this fixed gravitational background. Stochastic quantization is performed by introducing an extra fictitious time variable τ , and solving the corresponding Langevin equations. Thus, we can start by introducing an extra time variable for the fermionic fields $\psi(x) \rightarrow \psi(x, \tau)$, $\bar{\psi}(x) \rightarrow \bar{\psi}(x, \tau)$, and then writing down the Langevin equations [53]-[56]. As the action in these Langevin equations is a deformed Dirac action, we have to include an appropriate Kernel $K(x, y)$ to ensure the relaxation process is such that the systems will approach equilibrium as $\tau \rightarrow 0$. We also need to introduce $\eta(x, \tau)$ and $\bar{\eta}(x, \tau)$ as anticommuting fermionic Gaussian noise. Here $\eta(x, \tau)$ and $\bar{\eta}(x, \tau)$ satisfy

$$\begin{aligned}\langle \eta(x, \tau) \rangle &= 0 \\ \langle \bar{\eta}(x, \tau) \rangle &= 0, \\ \langle \eta(x', \tau') \bar{\eta}(x, \tau) \rangle &= 2\delta(\tau - \tau') K(x, x') \delta^4(x - x').\end{aligned}\quad (41)$$

Now using this anticommuting fermionic Gaussian noise, we can write the Langevin equations as

$$\begin{aligned}\frac{\partial \psi(x, \tau)}{\partial \tau} &= - \int d^4y K(x, y) \frac{\delta S[\psi, \bar{\psi}]}{\delta \bar{\psi}} + \eta(x, \tau), \\ \frac{\partial \bar{\psi}(x, \tau)}{\partial \tau} &= \int d^4y \frac{\delta S[\psi, \bar{\psi}]}{\delta \psi} K(x, y) + \bar{\eta}(x, \tau),\end{aligned}\quad (42)$$

where the action for the deformed Dirac equation on curved spacetime is given by

$$S[\psi, \bar{\psi}] = \int d^4x e\bar{\psi}[i\gamma^\mu (1 - \alpha\sqrt{-D^\nu D_\nu} - 2\alpha^2 D^\nu D_\nu) D_\mu - m]\psi. \quad (43)$$

As we are analysing the system semi-classically, our dynamical variables are only ψ and $\bar{\psi}$. The partition function for the deformed Dirac equation can be written as

$$Z = \int D\bar{\eta} D\eta \exp\left(-\frac{1}{2} \int d^4x d^4y d\tau \bar{\eta}(x, \tau) K^{-1}(x, y) \eta(y, \tau)\right). \quad (44)$$

After writing down the partition function, we can change $\bar{\eta}, \eta$ to $\bar{\psi}, \psi$, and thus obtain

$$\begin{aligned} Z &= \int D\bar{\psi} D\psi \det\left[\frac{\delta\bar{\eta}}{\delta\psi}\right]^{-1} \det\left[\frac{\delta\eta}{\delta\bar{\psi}}\right]^{-1} \\ &\quad \times \exp\left(-\frac{1}{2} \int d^4x d^4y d\tau \bar{\eta}(x, \tau) K^{-1}(x, y) \eta(y, \tau)\right). \end{aligned} \quad (45)$$

Now using Langevin equations, we obtain

$$\begin{aligned} Z &= \int D\bar{\psi} D\psi \det\left[K^{-1} \frac{\partial}{\partial\tau} - \frac{\delta^2 S[\bar{\psi}, \psi]}{\delta\bar{\psi}\delta\psi}\right]^{-1} \det\left[K^{-1} \frac{\partial}{\partial\tau} + \frac{\delta^2 S[\bar{\psi}, \psi]}{\delta\psi\delta\bar{\psi}}\right]^{-1} \\ &\quad \times \exp\left(-\frac{1}{2} \int d^4x d^4y d\tau \left[\frac{\partial\bar{\psi}}{\partial\tau} K^{-1} - \frac{\delta S[\bar{\psi}, \psi]}{\delta\psi}\right]\right. \\ &\quad \left.\times \left[\frac{\partial\psi}{\partial\tau} + K \frac{\delta S[\bar{\psi}, \psi]}{\delta\bar{\psi}}\right]\right), \end{aligned} \quad (46)$$

where the determinant can be defined as the regularized product of eigenvalues. Thus, we can obtain an expression for the determinant in terms of ghost fields $(c_1, \bar{c}_1, c_2, \bar{c}_2)$, because we can write the following expression [71],

$$\begin{aligned} \bar{c}_1 \det\left[K^{-1} \frac{\partial}{\partial\tau} - \frac{\delta^2 S}{\delta\bar{\psi}\delta\psi}\right] c_1 &= \lambda \bar{c}_1 c_1, \\ \bar{c}_2 \det\left[K^{-1} \frac{\partial}{\partial\tau} + \frac{\delta^2 S}{\delta\psi\delta\bar{\psi}}\right] c_2 &= \lambda \bar{c}_2 c_2. \end{aligned} \quad (47)$$

With this partition function, we can write

$$\begin{aligned} \mathcal{L}_{eff} &= \frac{1}{2} \frac{\partial\bar{\psi}}{\partial\tau} K^{-1} \frac{\partial\psi}{\partial\tau} - \frac{1}{2} \frac{\delta S[\bar{\psi}, \psi]}{\delta\bar{\psi}} K \frac{\delta S[\bar{\psi}, \psi]}{\delta\psi} \\ &\quad + \bar{c}_1 \left[K^{-1} \frac{\partial}{\partial\tau} - \frac{\delta^2 S[\bar{\psi}, \psi]}{\delta\bar{\psi}\delta\psi}\right] c_1 \\ &\quad + \bar{c}_2 \left[K^{-1} \frac{\partial}{\partial\tau} + \frac{\delta^2 S[\bar{\psi}, \psi]}{\delta\psi\delta\bar{\psi}}\right] c_2. \end{aligned} \quad (48)$$

We can introduce auxiliary fields $\bar{F}$ and F and write the partition function for this theory as [72]-[73]

$$Z = \int D\bar{\psi} D\psi Dc_1 D\bar{c}_1 Dc_2 D\bar{c}_2 DF D\bar{F} \exp\left(-\int d^4x e\mathcal{L}_{eff}\right), \quad (49)$$

where now

$$\begin{aligned}
\mathcal{L}_{eff} = & 2\bar{F}K^{-1}F + i \left[\frac{\partial\bar{\psi}}{\partial\tau}K^{-1} - \frac{\delta S[\bar{\psi}, \psi]}{\delta\psi} \right] F \\
& + i\bar{F} \left[\frac{\partial\psi}{\partial\tau} + K \frac{\delta S[\bar{\psi}, \psi]}{\delta\bar{\psi}} \right] + \bar{c}_1 \left[K^{-1} \frac{\partial}{\partial\tau} - \frac{\delta^2 S[\bar{\psi}, \psi]}{\delta\bar{\psi}\delta\psi} \right] c_1 \\
& + \bar{c}_2 \left[K^{-1} \frac{\partial}{\partial\tau} + \frac{\delta^2 S[\bar{\psi}, \psi]}{\delta\psi\delta\bar{\psi}} \right] c_2.
\end{aligned} \tag{50}$$

We can express this action using the superfield formalism. In order to do that, we define superfields $\Omega(x, \tau, \theta, \bar{\theta})$ and $\bar{\Omega}(x, \tau, \theta, \bar{\theta})$, as

$$\begin{aligned}
\Omega(x, \tau, \theta, \bar{\theta}) &= \psi(x, \tau) + \bar{\theta}c_1(x, \tau) + \bar{c}_2(x, \tau)\theta + i\theta\bar{\theta}F(x, \tau), \\
\bar{\Omega}(x, \tau, \theta, \bar{\theta}) &= \bar{\psi}(x, \tau) + \bar{\theta}c_2(x, \tau) + \bar{c}_1(x, \tau)\theta + i\theta\bar{\theta}\bar{F}(x, \tau).
\end{aligned} \tag{51}$$

We also define superderivatives and supercharges as

$$\begin{aligned}
D &= \frac{\partial}{\partial\theta} - \theta \frac{\partial}{\partial\tau}, & Q &= \frac{\partial}{\partial\theta}, \\
\bar{Q} &= \frac{\partial}{\partial\theta} + \bar{\theta} \frac{\partial}{\partial\tau}, & \bar{D} &= \frac{\partial}{\partial\theta},
\end{aligned} \tag{52}$$

where

$$\{D, \bar{D}\} = -\frac{\partial}{\partial\tau}, \quad \{Q, \bar{Q}\} = \frac{\partial}{\partial\tau}. \tag{53}$$

The action $S = S_1 + S_2 + S_3$ coincides with the component action, if we make the following changes $\bar{F}K^{-1} \rightarrow \bar{F}$, $\bar{\psi}K^{-1} \rightarrow \bar{\psi}$, $\bar{c}_1K^{-1} \rightarrow \bar{c}_1$, $\bar{c}_2K^{-1} \rightarrow \bar{c}_2$, and $F \rightarrow F$, $\psi \rightarrow \psi$, $c_1 \rightarrow c_1$, $c_2 \rightarrow c_2$ [76], where

$$\begin{aligned}
S_1 &= \int d^4x d\bar{\theta}d\theta \bar{D}\bar{\Omega}D\Omega \\
&= \int d^4x e \left[\bar{c}_1 \frac{\partial c_1}{\partial\tau} + \bar{F}F + i\bar{F} \frac{\partial\psi}{\partial\tau} \right], \\
S_2 &= \int d^4x d\bar{\theta}d\theta \bar{D}\Omega D\bar{\Omega} \\
&= \int d^4x e \left[\bar{c}_2 \frac{\partial c_2}{\partial\tau} + F\bar{F} + i\bar{F} \frac{\partial\bar{\psi}}{\partial\tau} \right], \\
S_3 &= \int d^4x d\bar{\theta}d\theta S[\bar{\Omega}, \Omega] \\
&= \int d^4x e \left[\bar{F} \frac{\delta S_3}{\delta\bar{\psi}} + \frac{\delta S_3}{\delta\psi} F + \bar{c}_1 \frac{\delta^2 S_3}{\delta\psi\delta\bar{\psi}} c_1 + \bar{c}_2 \frac{\delta^2 S_3}{\delta\bar{\psi}\delta\psi} c_2 \right].
\end{aligned} \tag{54}$$

Thus, stochastic quantization of the deformed Dirac equation can be done using superspace formalism. It may be noted that we have performed the stochastic quantization of the deformed Dirac equation using the semi-classical approximation. Thus, during this analysis, we have taken the gravitational field as a fixed background field, and performed the stochastic quantization of the deformed Dirac equation on this fixed background field.

6 Conclusion

In this paper, we first clarified what it means to have a minimum measurable time in the theory. This was done by first defining the observable time as the time of occurrence of an certain event at which some specified value gets assigned to a dynamical quantity [45]-[50]. Then viewing minimum measurable time as the existence of a minimum measurable value to the certainty with which the occurrence of that event can be measured by the appearance of this specified value for that dynamical quantity. Then we have combined the deformation of the Heisenberg algebra caused by the doubly special relativity and with its deformation caused by the existence of minimum measurable length and time scales. We also have completed this deformed Heisenberg algebra by including a covariant deformation of its temporal part. It may be noted that one of the most interesting consequence of the spatial part of this deformed Heisenberg algebra was that it leads to the discretization of space [35]. It will be interesting to repeat this analysis for the full Heisenberg algebra. It is expected that such an analysis will lead to the discretization of time. It was observed that this deformed algebra gives rise to non-local fractional derivative terms. However, a formal meaning was given to these non-local terms in the framework of harmonic extension of functions. This deformed Dirac equation was also analysed on curved spacetime. It was demonstrated that this deformation of the Dirac equation also deforms the geometry of spacetime. Thus, corrections to the Einsteins equations were obtained by analysing this deformation. Finally, the stochastic quantization of this deformed Dirac equation was performed on curved spacetime. It was done by using a fictitious time coordinate and anticommuting fermionic Gaussian noise. Thus, we were able to analyse the Langevin equation for the deformed Dirac action on curved spacetime. It was also observed that the stochastic quantization of this deformed Dirac equations could be done using superspace formalism. It may be noted that we did not analyse the stochastic quantization of gravitational field in this model. It would be interesting to analyse the stochastic quantization of this modified theory of gravity in the framework of perturbative quantum gravity. However, as not all degrees of freedom in this modified theory of gravity are physical, we will need to add gauge fixing and ghost terms, to the original action. This resultant effective action, which will be given by a sum of the original action with the gauge fixing and ghost terms, will be invariant under a new symmetry called the BRST symmetry. In fact, various consequences of the BRST symmetry of perturbative quantum gravity have been analysed [77]-[82]. It would be interesting to analyse these consequences of the BRST symmetry for this modified theory of gravity.

Non-local fractional derivative terms also occur in fermionic theories with Lifshitz scaling, and stochastic quantization of such Lifshitz theories has also been performed [76]. It would thus be interesting to combine the effects coming from the deformation of the Dirac equation with Lifshitz scaling. We expect that non-local fractional derivative terms, that will occur due to the combination of the deformation of the Heisenberg algebra with Lifshitz scaling, could still be analysed using the harmonic extension of functions. It is also expected that by analysing the deformed Dirac equation with Lifshitz scaling, further modification to the geometry will occur. It would be interesting to analyse this modification to the Einstein equations. Corrections to various cosmological

models can be calculated from these modified Einstein equations. Furthermore, phenomenological predications about cosmology can also be made from such corrected cosmological models. It may be noted that the deformation of the Wheeler-DeWitt equation caused by this algebra has already been analysed [44]. It may be noted that there are various proposals for the problem of time in quantum gravity [83]-[84], and it would be interesting to analyse the deformation of models based on such proposals. Thus, it would also be interesting to analyse the modification to quantum cosmology that occurs because of this deformed algebra. It may also be noted that an alternative deformation of the Heisenberg algebra can lead to noncommutative spacetime [85]-[86]. The noncommutativity has been generalized to non-anticommutativity, and perturbative quantum gravity on non-anticommutative spacetime has been analysed [87]-[89]. It would be interesting to combine the deformation of the Heisenberg algebra studied in this paper with non-anticommutativity, and analyse the effect of this combined deformation on quantum gravity. Recently, it has also been observed that all black objects in gravity's rainbow will have a remnant [90], and this has phenomenological consequences for detection of higher dimensional black holes at the LHC [91]. As both gravity's rainbow and the modification of gravity studied in this paper are motivated by doubly special relativity, it would be interesting to analyse black objects in the modified theory of gravity discussed in this paper. It is possible that a similar result may hold for all black objects in this modified theory of gravity.

References

- [1] D. Amati, M. Ciafaloni and G. Veneziano, Phys. Lett. B 216, 41 (1989)
- [2] A. Kempf, G. Mangano, and R. B. Mann, Phys. Rev. D 52, 1108 (1995)
- [3] L. N. Chang, D. Minic, N. Okamura, and T. Takeuchi, Phys.Rev. D65, 125027 (2002)
- [4] L. N. Chang, D. Minic, N. Okamura, and T. Takeuchi, Phys. Rev. D65, 125028 (2002)
- [5] S. Benczik, L. N. Chang, D. Minic, N. Okamura, S. Rayyan, and T. Takeuchi, Phys. Rev. D66, 026003 (2002)
- [6] P. Dzierzak, J. Jezierski, P. Malkiewicz, and W. Piechocki, Acta Phys. Polon. B41, 717 (2010)
- [7] M. Maggiore, Phys. Lett. B304, 65 (1993)
- [8] M. I. Park, Phys. Lett. B659, 698 (2008)
- [9] D. Amati, M. Ciafaloni, and G. Veneziano, Phys. Lett. B 216, 41 (1989)
- [10] M. Maggiore, Phys. Lett. B 304, 65 (1993)
- [11] M. Maggiore, Phys. Rev. D 49, 5182 (1994)
- [12] M. Maggiore, Phys. Lett. B 319, 83 (1993)
- [13] L. J. Garay, Int. J. Mod. Phys. A 10, 145 (1995)

- [14] F. Scardigli, Phys. Lett. B 452, 39 (1999)
- [15] C. Bambi, F. R. Urban, Class. Quantum Grav. 25, 095006 (2008)
- [16] A. Kempf, G. Mangano, and R. B. Mann, Phys. Rev. D 52, 1108 (1995)
- [17] A. Kempf, J. Phys. A 30, 2093 (1997)
- [18] F. Brau, J. Phys. A 32, 7691 (1999)
- [19] K. Nozari, and B. Fazlpour, Chaos, Solitons and Fractals, 34, 224 (2007)
- [20] K. Nozari, Phys. Lett. B. 629, 41 (2005)
- [21] K. Nozari, T. Azizi, Gen. Relativ. Gravit. 38, 735 (2006)
- [22] P. Pedram, Int. J. Mod. Phys. D 19, 2003 (2010)
- [23] S. Das, and E. C. Vagenas, Phys. Rev. Lett. 101, 221301 (2008)
- [24] J. Magueijo, and L. Smolin, Phys. Rev. Lett. 88, 190403 (2002)
- [25] J. Magueijo, and L. Smolin, Phys. Rev. D 71, 026010 (2005)
- [26] J. L. Cortes and J. Gamboa, Phys. Rev. D 71, 065015 (2005)
- [27] G. 't Hooft, Class. Quant. Grav. 13, 1023 (1996).
- [28] V. A. Kostelecky and S. Samuel, Phys. Rev. D 39, 683 (1989)
- [29] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and S. Sarkar, Nature 393, 763 (1998).
- [30] R. Gambini and J. Pullin, Phys. Rev. D 59, 124021 (1999).
- [31] S. M. Carroll, J. A. Harvey, V. A. Kostelecky, C. D. Lane and T. Okamoto, Phys. Rev. Lett. 87, 141601 (2001).
- [32] P. Horava, Phys. Rev. D 79, 084008 (2009).
- [33] J. Magueijo and L. Smolin, Class. Quant. Grav. 21, 1725 (2004)
- [34] J. J. Peng, and S. Q. Wu, Gen. Rel. Grav. 40, 2619 (2008)
- [35] A. F. Ali, S. Das and E. C. Vagenas, Phys. Lett. B 678, 497 (2009)
- [36] S. Das, E. C. Vagenas, and A. F. Ali, Phys. Lett. B 690, 407 (2010)
- [37] P. Pedram, Europhys. Lett. 89, 50008 (2010)
- [38] P. Pedram, K. Nozari, and S. H. Taheri, JHEP. 1103, 093 (2011)
- [39] A. F. Ali, S. Das, and E. C. Vagenas, Phys. Rev. D84, 044013 (2011)
- [40] V. Husain, D. Kothawala and S. S. Seahra, Phys. Rev. D 87, 025014 (2013)
- [41] M. Kober, Phys. Rev. D82, 085017 (2010)
- [42] M. Kober, Int. J. Mod. Phys. A 26, 4251 (2011)

- [43] M. Faizal, Int. J. Geom. Meth. Mod. Phys. 12, 1550022 (2015)
- [44] M. Faizal, Int. J. Mod. Phys. A 29, 1450106 (2014)
- [45] P. Buscha, M. Grabowskib and P. J. Lahtic, Phys. Lett. A 191, 357 (1994)
- [46] R. S. Raghavan, D. Minic, T. Takeuchi and C. H. Tze, arXiv:1210.5639
- [47] V. S. Olkhovsky, Adv. Math. Phys. 2009, 859710 (2009)
- [48] V. S. Olkhovsky, Int. J. Mod. Phys. A22, 5063 (2007)
- [49] V. S. Olkhovsky and E. Recami, Int. J. Mod. Phys. B 22, 1877 (2008)
- [50] R. Brunetti, K. Fredenhagen and M. Hoge, Found. Phys. 40, 1368 (2010)
- [51] H. Nikolic, Phys. Lett. B 678, 218 (2009)
- [52] H. Nikolic, arXiv:1407.8058
- [53] G. Parisi and Y. S. Wu, Sci. Sin. 24, 483 (1981)
- [54] A. Barchielli and G. Lupieri, J. Math. Phys. 26, 2222 (1985)
- [55] F. Guerra and R. Marra, Phys. Rev. D29, 1647 (1984)
- [56] P. H. Damgaard and K. Tsokos, Nucl. Phys. B 235, 75 (1983)
- [57] R. A. Janik, M. A. Nowak, G. Papp and I. Zahed, Phys. Rev. Lett. 81, 264 (1998)
- [58] J. C. Osborn and J. J. M. Verbaarschot, Phys. Rev. Lett. 81, 268 (1998)
- [59] J. C. Osborn and J. J. M. Verbaarschot, Phys. Rev. Lett. 81, 268 (1998)
- [60] T. Guhr, T. Wilke and H.A. Weidenmuller, Phys. Rev. Lett. 85, 2252 (2000)
- [61] Y. M. Cho, Phys. Rev. D 14, 2521 (1976)
- [62] J. W. Maluf, Phys. Rev. D 67, 108501 (2003)
- [63] F. W. Hehl, P. Von Der Heyde, G. D. Kerlick and J. M. Nester, Rev. Mod. Phys. 48, 393 (1976)
- [64] I. Piovski, M. R. Vanner, M. Aspelmeyer, M. Kim and C. Brukner, Nature Phys. 8, 393 (2012)
- [65] R. T. Seeley, Proc. Symp. Pure Math. 10, 288 (1967)
- [66] L. Caffarelli, and L. Silvestre, Comm. Part. Diff. Eqs. 32, 1245 (2007)
- [67] J. J. Giambiagi, Nuovo Cim. A 104, 1841 (1991)
- [68] C. G. Bollini and J. J. Giambiagi, J. Math. Phys. 34, 610 (1993)
- [69] C. Laemmerzahl, J. Math. Phys. 34, 3918 (1993)
- [70] J. Tan, Calc. Var. 42, 21 (2011)

- [71] U. Lindstrom, M. Rocek, W. Siegel, P. van Nieuwenhuizen, and A. E. van de Ven, J. Math. Phys. 31, 1761 (1990)
- [72] G. Parisi and N. Sourlas, Nucl. Phys. B 206, 321 (1982)
- [73] S. Chaturvedi, A. K. Kapoor and V. Srinivasan, Phys. Lett. B 140, 56 (1984)
- [74] A. I. Ahieser and V. B. Berestetskii, Quantum Electrodynamics. New York: Wiley Interscience, (1969)
- [75] S. I. Kruglov, Symmetry and Electromagnetic Interaction of Fields with Multi-Spin. Nova Science Publishers, Huntington, New York, (2001)
- [76] H. Montani and F. A. Schaposnik, Phys. Rev. D 86, 065024 (2012)
- [77] N. Nakanishi and I. Ojima, Covariant operator formalism of gauge theories and quantum gravity, World Sci. Lect. Notes. Phys. (1990)
- [78] M. Faizal, Found. Phys. 41, 270 (2011)
- [79] M. Faizal, J. Phys. A 44, 402001 (2011)
- [80] Kugo and I. Ojima, Nucl. Phys. B 144, 234 (1978)
- [81] M. Faizal, Mod. Phys. Lett. A 28, 1350034 (2013)
- [82] M. Mena and M. Tahiri, Phys. Rev. D 57, 7312 (1998)
- [83] Barbour, Phys. Rev. D 47, 5422 (1993)
- [84] G. W. Gibbons, Fundam. Theor. Phys. 172, 109 (2012)
- [85] S. Minwalla, M. V. Raamsdonk and N. Seiberg, JHEP. 0002, 020 (2000)
- [86] N. Seiberg and E. Witten, JHEP. 9909, 032 (1999)
- [87] M. Faizal, Mod. Phys. Lett. A 27, 1250075 (2012)
- [88] M. Faizal, Phys. Lett. B 705, 120 (2011)
- [89] J. W. Moffat, Phys. Lett. B 491, 345 (2000)
- [90] A. F. Ali, M. Faizal and Mohammed M. Khalil, Nucl. Phys. B 894, 341 (2015)
- [91] A. F. Ali, M. Faizal and Mohammed M. Khalil, Phys. Lett. B 743, 295 (2015)